

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed and, completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302 and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE April 17, 2001		3. REPORT TYPE AND DATES COVERED Technical Report
4. TITLE AND SUBTITLE Syntheses of Diazadithiacrown Ethers Containing Two 8-Hydroxyquinoline Side Arms			5. FUNDING NUMBERS N00014-98-1-0485 98PR05020-00	
6. AUTHOR(S) H.-C. Song, J. S. Bradshaw, Y.-W. Chen, G.-P. Xue, J. A. Chiara, K. E. Krakowiak, P. B. Savage, Z.-L. Xue, and R. M. Izatt				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry and Biochemistry Brigham Young University Provo, UT 84602			8. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No 27	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Dr. Stephen W. McElvany (Program Officer) Office of Naval Research 800 No Quincy Street Arlington, VA 22217-5660			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Ten new diazadithiacrown ethers containing two 8-hydroxyquinoline (HQ) sidearms attached through the HQ 7-positions and four new diazadithiacrown ethers containing two HQ sidearms attached through the HQ 2-positions have been prepared. Some of these new ligands also contain a hydroxymethyl substituent. The starting macrocyclic diazadithiacrown ethers were obtained by treatment of a bis(-chloroamide) with the appropriate dimercaptan using K_2CO_3 as the base followed by reduction of the resulting macrocyclic dithiadamide by BH_3-THF or by $NaBH_4$ in the presence of BF_3-ether as a catalyst. HQ-containing ligands 23-32 were synthesized by a Mannich reaction of the secondary macrocyclic diamines with the substituted-8-hydroxyquinoline. HQ-containing ligands 33-36 were prepared by reductive amination of the secondary macrocyclic diamines with 8-hydroxyquinoline-2-carbaldehyde.</p>				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
20010501 090			16. PRICE CODE NA	
			20. LIMITATION OF ABSTRACT Unlimited	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified		

OFFICE OF NAVAL RESEARCH

Grant N00014-98-1-0485

Technical Report No. 27

**Syntheses of Diazadithiacrown Ethers Containing Two
8-Hydroxyquinoline Side Arms**

by

H.-C. Song, J. S. Bradshaw, Y.-W. Chen, G.-P. Xue, J. A. Chiara,
K. E. Krakowiak, P. B. Savage, Z.-L. Xue, and R. M. Izatt

Department of Chemistry and Biochemistry
Brigham Young University, Provo, UT 84602-4678

April 17, 2001

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

Syntheses of Diazadithiacrown Ethers Containing Two 8-Hydroxyquinoline Side Arms

Hua-Can Song,^a Jerald S. Bradshaw,^b Yi-Wen Chen,^a Guo-Ping Xue,^b Joseph A. Chiara,^b Krzysztof E. Krakowiak,^b Paul B. Savage,^b Zun-Le Xue,^a and Reed. M. Izatt^b

^aDepartment of Chemistry, Zhongshan University, Guangzhou 510275 P. R. China

^bDepartment of Chemistry & Biochemistry, BYU, Provo, UT 84602, USA

Introduction

In general, the complexing ability and selectivity of lariat ethers for metal ions can be varied by changing certain parameters such as the acidity of the phenolic OH group; the size of the crown ether ring; type, number, and position of the complexing crown ether heteroatoms; the stereochemistry imposed by the arms which connect the phenolic group to the macroring; and the pH of media. For example, diaza-18-crown-6 containing two 5-chloro-8-hydroxyquinoline (CHQ) groups attached through the CHQ 7-position (**1**, Figure 1) exhibits a stronger affinity for Mg^{2+} than for Ba^{2+} (log K value in MeOH for Mg^{2+} is 6.82, for Ba^{2+} 3.60) and its isomer, diaza-18-crown-6 bearing two CHQ groups attached through the CHQ 2-position (**4**), has a stronger affinity for Ba^{2+} than for Mg^{2+} (log K value in MeOH for Ba^{2+} is 12.2).¹ Ligands **3** and **5**, the HQ analogs of **1** and **4**, respectively (**1** and **4** with the chlorine atoms removed), do not exhibit the same complexing properties as do **1** and **4**.² Increasing the number of macroring nitrogen atoms and changing the size of macroring could change the affinity of the ligand toward the heavy metal ions. For example, for ligand **5**, the log K value in MeOH for Cu^{2+} is 4.39 while its tetraaza-15-crown-5 analog (**6**) has a log K value for Cu^{2+} of 15.5.³ Ligand **2**, which has a 5-nitro substituent on each 8-hydroxyquinoline, has a high affinity and selectivity for Hg^{2+} and has proven to be a chemosensor for Hg^{2+} .⁴ Diaza-18-crown-6 with two 4-methyl(or nitro)-6-aminophenol groups attached through the phenol 2-positions (**7** and **8**) form dinuclear complexes with one Cu^{2+} complexed to the two 6-aminophenols and one Na^+ in the macroring cavity.⁵

Diazadithia(or trithia)crown ethers **9** containing two HQ side arms have also been synthesized.^{6,7} These new azathia ligands have poor solubilities in MeOH and, therefore, their complexing properties with metal ions cannot be conveniently studied. A few of ligands **9** have a hydroxymethyl substituent attached to the macroring and are thereby more soluble in methanol. Herein, we report the synthesis of a series of new diazadithiacrown ethers bearing two 5-substituent(or 2-methyl)-HQ side-arms. Some of these new ligands contain a hydroxymethyl group on the macroring. A report on the affinities of some of these new ligands for metal ions and their possible use as sensors for metal ions will be reported in due course.

Results and Discussion

The CHQ and HQ side arms are best attached to the diazadithiacrown ethers through macroring NH groups. Macrocyclic ligands containing two secondary amine functions have conveniently been prepared by the crab-like synthesis using bis(α -chloroamide)s.^{3,6,8-10} The NH functions of the secondary bis(α -chloroamide)s are unreactive toward alkylating agents including thiols.^{6,11} In the present case, bis(α -chloroamide)s **10-12** were treated with the appropriate dimercaptans using K_2CO_3 as the base to form macrocyclic diamides **13-15** in yields of 46%-61% as shown in Scheme 1. The macrocyclic diamides were in turn reduced to the desired diazadithiacrown ethers **16, 17, and 22** by either B_2H_6 -THF or the $NaBH_4$ - BF_3 -THF complex (Scheme 1). Ligands **18-21** shown in Scheme 1 were prepared as reported.⁶ Satisfactory elemental analyses were obtained for the new macrocyclic diamides or for new HQ and CHQ armed ligands **23-36** prepared from them.

The Mannich aminomethylation reaction has been used to attach HQ and CHQ groups to the azacrown ethers through HQ and CHQ 7-positions.^{2,6,12-14} In the present case, the appropriate diazadithiacrown ether (**16-22**) and the appropriate HQ derivative were treated with paraformaldehyde in refluxing benzene in the one-step aminomethylation reaction^{2,6,15,16} to give the bis(2- or 5-substituted-8-quinolin-7-ylmethyl)-substituted ligands **23-32** (Scheme 2).

The products of the Mannich reaction of the diazadithiacrown ethers with HQ (**23 and 27**) and 8-hydroxyquinaldine (**26**) were mixtures. Each of these two starting materials has no substituent on the quinoline 5-position. Thus, both the 5 and 7 positions could be aminomethylated under these reaction conditions. Although we did not look for the side products in these reactions, we recently showed by a careful 1H NMR analysis that when diazatrithiacrown ether **18** was treated with 8-hydroxyquinaldine, the product mixture proved to be about 90% of the desired product where both quinoline substituents were attached through the 8-hydroxyquinaldine 7-position, about 9% of the product with one 8-hydroxyquinaldine attached through its 7-position and the other through its 5-position and the remaining product had both 8-hydroxyquinaldine groups attached through their 5-positions. Thus, we suspect that products **23, 26 and 27** are mixtures where the HQ groups are attached through their 7- and 5-positions.

HQ has been attached to diaza-18-crown-6² and a series of tetraaza-15 (and 16)-crown-5 ligands³ through the HQ 2-position by a reductive amination process using $NaBH(OAc)_3$.¹⁷ In the present case, 8-hydroxyquinoline-2-carbaldehyde and the appropriate ligand (**16-18 or 20**) were treated with $NaBH(OAc)_3$ to form the bis(8-hydroxyquinolin-2-ylmethyl)-substituted ligands **33-36** in yields of 46% - 66% (Scheme 3). It is important to note that the hydroxy group of HQ did not have to be protected for this reaction as previously reported.³







Figure 1. Compounds Mentioned in the Introduction.



Scheme 1. Preparation of Diazadithiacrown Ethers **16**, **17** and **22**.



Scheme 2. Preparation of 8-Hydroxyquinoline-substituted Diazadithiacrown Ethers **23-32** by the Mannich Reaction.



Scheme 3. Preparation of 8-Hydroxyquinoline-substituted Diazadithiacrown Ethers **33-36** by Reductive Amination.

References

1. Zhang, X.X.; Bordunov, A.V.; Bradshaw, J.S.; Dalley, N.K.; Kou, X.; Izatt, R.M. *J. Am. Chem. Soc.* **1995**, *117*, 11507.
2. Su, N.; Bradshaw, J.S.; Zhang, X.X.; Song, H.-C.; Xue, G.-P.; Savage, P.B.; Krakowiak, K.E.; Izatt, R.M. *J. Org. Chem.* **1999**, *64*, 8855.
3. Yang, Z.; Bradshaw, J.S.; Zhang, X.X.; Savage, P.B.; Krakowiak, K.E.; Dalley, N.K.; Su, N.; Bronson, R.T.; Izatt, R.M. *J. Org. Chem.* **1999**, *64*, 3162.
4. Prodi, L.; Bargossi, C.; Montalti, M.; Zaccheroni, N.; Su, N.; Bradshaw J.S.; Izatt, R.M.; Savage, P.B. *J. Am. Chem. Soc.* **2000**, *122*, 6769.
5. Su, N.; Bradshaw, J.S.; Zhang, X.X.; Savage, P.B.; Krakowiak, K.E.; Izatt, R.M.; *J. Org. Chem.* **1999**, *64*, 3825.
6. Bradshaw J. S.; Song H-C; Xue G-P, Bronson T. R.; Chiara J. A.; Krakowiak K. E.; Savage P. B.; Izatt R. M. *Supramol. Chem.* in press.
7. Xue G-P; Bradshaw J. S.; Song H-C; Bronson T. R.; Savage P. B.; Krakowiak K. E.; Izatt R. M.; Prodi L.; Montalti M.; Zaccheroni N. *Tetrahedron* **2001**, *57*, 87.
8. Bradshaw, J.S.; Krakowiak, K.E.; Izatt, R.M. *J. Heterocyclic Chem.* **1989**, *26*, 1431.
9. Krakowiak, K.E.; Bradshaw, J.S.; Izatt, R.M. *J. Org. Chem.* **1990**, *55*, 3364.
10. Krakowiak, K.E.; Bradshaw, J.S.; Izatt, R.M. *Synlett.* **1993**, 611.
11. Bradshaw, J.S.; Krakowiak, K.E.; An, H.-Y.; Izatt, R.M. *J. Heterocyclic Chem.* **1990**, *27*, 2113.
12. Lukyanenko, N.K.; Pastushok, A.N.; Bordunov, A.V.; Vetrogon, V.I.; Vetrogon, N.I.; Bradshaw, J.S. *J. Chem. Soc. Perkin Trans. 1*, **1994**, 1489.
13. Bordunov, A.V.; Bradshaw, J.S.; Zhang, X.X.; Dalley, N.K.; Kou, X.; Izatt, R.M. *Inorg. Chem.* **1996**, *35*, 7229.
14. Bordunov, A.V.; Bradshaw, J.S.; Pastushok, V.N; Izatt, R.M. *Synlett.*, **1996**, 933
15. Chi, K.-W.; Wei, H.-C., Kottke, T.; Lagow, R.J. *J. Org. Chem.* **1996**, *61*, 5684.
16. Su, N.; Bradshaw, J.S.; Savage, P.B.; Krakowiak, K.E.; Izatt, R.M.; DeWall, S.L.; Gokel, G.W. *Tetrahedron* **1999**, *55*, 9737.
17. Abdel-Magid, A.F.; Carson, K.G.; Harris, B.D.; Maryanoff, C.A.; Shah, R.D. *J. Org. Chem.* **1996**, *61*, 3849.